

Proceedings and Abstracts of the 15<sup>th</sup> Annual Scientific Meeting of the Society for Healthcare Epidemiology of America; 2005 April 9-12; Los Angeles, California, United States of America; 2005. p. 134.

The Attributable Cost, And Length Of Hospital Stay Of Central Line Associated Blood Stream Infection In Intensive Care Units In Mexico. A Prospective, Matched Analysis

Francisco Higuera, MD<sup>1</sup>, MS Rangel-Frausto, MD<sup>2</sup>, Victor D. Rosenthal, MD, CIC, MSc<sup>3</sup>, Jorge Castanon, MD<sup>2</sup>, Guillermo Franco, MD<sup>1</sup>, Natividad Tobal, RN<sup>2</sup>, Pablo Duarte, MD<sup>4</sup>, J Martinez Soto, MD<sup>5</sup>, Javier Ruiz, MD<sup>1</sup>.

<sup>1</sup>General Hospital, Mexico, Mexico, <sup>2</sup>IMSS, Mexico, Mexico, <sup>3</sup>Medical College of Buenos Aires, Buenos Aires, Argentina, <sup>4</sup>General Hospital, Mexico, Mexico, <sup>5</sup>IMSS Gabriel Mancera, Mexico, Mexico.

Background:

Limited information is available on the financial impact of central venous catheter (CVC)-associated bloodstream infection (BSI) in Mexico.

Objectives:

In order to calculate the cost of BSI in intensive care units, a 2 - year prospective nested case-control study was undertaken at three hospitals in Mexico.

Methods:

This study was performed in four adult intensive care units (ICUs) from 3 Mexican hospitals. 55 patients with BSI (cases) and 55 patients without BSI (controls) were matched for hospital, ICU, timing of admission, length of stay, gender, age, and average severity of illness score. Patient's length of stay (LOS) in the ICU was obtained prospectively on daily rounds; the estimated cost of each ICU day was provided by the hospital's finance department; the average estimated cost of antibiotics prescribed for BSIs was provided by the hospital's pharmacy department.

Results:

The mean LOS of patients with BSI was 13.44 days. The mean LOS of patients without BSI was 7.38. The mean extra LOS for cases (compared to the controls) was 6.05 days.

The mean cost of patients with BSI was US\$ 28,728.71. The mean cost of patients without BSI was US\$ 21,860.35. The mean extra cost for cases (compared to the controls) was US\$ 6,899.11.

Conclusions:

CVC-associated BSI results in significant patient morbidity and is associated with considerable increased utilization of health care resources. In the present study, patients with BSI had significant prolongation of hospitalization, and cost. The present study supports the need for routine utilization of preventative interventions to reduce CVC-associated BSIs as a means of reducing health care costs.